

TRACE ELEMENT ABUNDANCES IN ST. AUBIN (UNGR IRON) GIANT CHROMITE AND ASSOCIATED PHASES.

G. Kurat¹, E. Zinner² and M. E. Varela³. ¹Naturhistorisches Museum, Vienna, Austria. gero.kurat@univie.ac.at. ²Phys. Dept., Washington University, St. Louis, MO, USA. ³Dept. Geol., UNS-CONICET, B. Blanca, BA, Argentina.

Introduction: Chromite crystals of up to 3 cm in size were recently described from the fine octahedrite St. Aubin [1]. The crystals are euhedral exhibiting triangular and hexagonal cross sections and are incompletely covered by schreibersite, troilite and swathing kamacite. They are accompanied by schreibersite, troilite, hibbingite and euhedral Fe-phosphate (sarcopsite or graffonite). We have analyzed the non-metallic phases for their major, minor and trace element contents with an EMP and an IMS-3f ion probe, which was also utilized to search for extinct ⁵³Mn in the Fe-phosphates – all following routine procedures.

Results: Chromite is pure FeCr₂O₄ containing (in ppm) ~8000 V, ~4700 Mn (but only ~600 Mg and ~0.4 Al), 0.15 Nb, 0.02 Sc and <0.003 Ce. The Fe-phosphate is also pure (Fe,Mn)₃(PO₄)₃ with (in ppm) ~18000 Mn, ~2000 Mg, ~270 Zn, 12.5 Cr, ~6 Co, 4.2 Ni, 0.01 Nb, 0.0017 Sc and ~0.0003 Ce. Hibbingite, Fe₂(OH)₃Cl (~18 wt% Cl), contains (in ppm) 6200 Ni, 2500 Co, 292 Mn, 0.0003 Sc and 0.0005 Ce. Fe-phosphates have excesses in ⁵³Cr with an initial ratio of ⁵³Mn/⁵⁵Mn = (1.5+/-0.3)x10⁻⁶.

Discussion: All non-metallic phases in St. Aubin are extremely poor in lithophile elements. Particularly striking are the very low contents of Al, Mg, Sc and Ti in chromite as compared to those reported by, e.g., [2]. Also, the contents of Zr and the REE are very low, all <0.01xCl. We interpret this to indicate derivation of the chromite from an environment that was very poor in all these elements. The same holds for the Fe-phosphate. Chromite and phosphate are also very poor in Ni, less so in the less siderophile Co, indicating equilibration with metal. On the other hand, chromite and phosphate are enriched in V and Nb and also Zn and Mn with respect to the common lithophile elements, indicating an elevated siderophile behavior of V and Nb – as was predicted by [3] - and formation of chromite and phosphate from reduced precursor phases. Hibbingite likely is a secondary phase after lawrencite and indicates also low abundances of lithophile elements during formation of the latter in the presence of metal.

Conclusion: All non-metallic phases in St. Aubin indicate formation in an environment that was very poor in lithophile elements. The abundance anomalies in V and Nb indicate reduced precursor phases (metals, carbides, etc.), which subsequently were oxidized to form chromite and Fe-phosphate early in the history, about contemporaneously with the Mn-Fe metasomatism event experienced by the angrites [4], while some ⁵³Mn still was alive. Thereafter, troilite and schreibersite formed, followed by lawrencite and final metal which preserved all phases.

Acknowledgements: We thank Andrey Gurenko and Nora Groschopf, Mainz, for permission to use and help with the EMPA, respectively; financial support was received in Austria from FWF, the Academy of Sciences and the Freunde des NHM in Wien, in Argentina from CONICET, and in the USA from NASA.

References: [1] Fehr and Carion 2004. MAPS 39, A139-A141. [2] Bunch et al. 1970. Contr. Mineral. Petrol. 25, 297-340. [3] Jurewicz et al. 1995. EPSL 132, 183-198. [4] Lugmair and Shukolyukov 2002. MAPS 37, 1001-1013.